

Some Results on R-KKM Mappings and R-KKM Selections and Their Applications

Ram U. Verma

metadata, citation and similar papers at core.ac.uk

Submitted by Ravi P. Agarwal

Received July 30, 1998

First some intersection theorems based on R-KKM mappings are obtained and then their application to minimax inequalities is given. © 1999 Academic Press

1. INTRODUCTION

The KKM theorem [5] and its generalizations turned out to be a powerful tool dealing with several nonlinear problems. Horvath [2] generalized the KKM theorem replacing the usual convexity assumptions by the topological notion of the contractibility. Motivated by the work of Horvath, Bardaro and Ceppitelli [1] generalized minimax inequalities for functions taking values in ordered vector spaces. At the same time, based on the notion of a convex hull operation introduced by Komiya [6], Joo' [3] and Joo' and Kassay [4] applied the concept of the pseudoconvexity obtaining some intersection theorems with their applications to minimax inequalities. Recently, the author [8–13] obtained some intersection theorems and their applications to minimax inequalities in generalized H-spaces. The aim of this paper is first to obtain an intersection theorem involving R-KKM selections, second to apply it to establish another intersection theorem involving R-KKM mappings, and finally, as a result, to derive some minimax inequalities. The obtained results generalize some results [1, 2, 7–12].

Let X be a topological space without linear structure, $P(X)$ denote the power set of X , and $\langle X \rangle$ the family of all nonempty finite subsets of X . Let Δ^n denote a standard $(n - 1)$ simplex $(e_1, \dots, e_2, \dots, e_n)$ in R^n .

Joo' [3] introduced, based on the notion of a convex space initiated by Komiya [6], the concept of a pseudoconvex space.



DEFINITION 1.1. A triple $(X, h, \{p\})$ is called a pseudoconvex space if:

(i) X is a topological space and $h: P(X) \rightarrow P(X)$ is a convex hull operation satisfying the properties: $h(\emptyset) = \emptyset$, $h(\{x\}) = \{x\}$ for all $x \in X$, $h(A) = \bigcup \{h(F): F \in \langle A \rangle\}$ for $A \subset X$, and $h(h(A)) = h(A)$ for $A \subset X$.

(ii) $p: \Delta^n \rightarrow h(F)$ is a continuous mapping from Δ^n onto $h(F)$, where Δ^n is a standard simplex of R^n and $F \in \langle X \rangle$.

(iii) For each $F \in \langle X \rangle$, p is convex hull preserving in the sense that $p(e_{i_1}, e_{i_2}, \dots, e_{i_k}) = h(\{x_{i_1}, x_{i_2}, \dots, x_{i_k}\})$ for all subsimplex $(e_{i_1}, e_{i_2}, \dots, e_{i_k}) \subset (e_1, e_2, \dots, e_n)$ and a subset $\{x_{i_1}, x_{i_2}, \dots, x_{i_k}\}$ of $\{x_1, x_2, \dots, x_n\}$ for $\{i_1, i_2, \dots, i_k\} \subset \{1, 2, \dots, n\}$.

A subset $K \subset X$ is called h -convex if $h(K) = K$.

DEFINITION 1.2. A triple $(X, H, \{p\})$ is said to be a generalized H -space (G - H -space) [8] if X is a topological space and $H: \langle X \rangle \rightarrow P(X) \setminus \{\emptyset\}$ is a mapping such that the following assumptions hold:

(i) For each $F, G \in \langle X \rangle$, there exists $F_1 \subset F$ such that $F_1 \subset G$ implies $H(F_1) \subset H(G)$.

(ii) For $F = \{x_1, x_2, \dots, x_n\} \in \langle X \rangle$, there exists a continuous mapping $p: \Delta^n \rightarrow H(F)$ such that for $\{i_1, i_2, \dots, i_k\} \subset \{1, 2, \dots, n\}$, we have

$$p(e_{i_1}, e_{i_2}, \dots, e_{i_k}) \subset H(\{x_{i_1}, x_{i_2}, \dots, x_{i_k}\}),$$

where $\{x_{i_1}, x_{i_2}, \dots, x_{i_k}\} \subset F$.

A subset K of X is called generalized H -convex (G - H -convex) if for each $F \in \langle X \rangle$ there exists a subset F_1 of F such that $F_1 \subset K$ implies $H(F_1) \subset K$.

DEFINITION 1.3. Let $(X, H, \{p\})$ be a G - H -space and $T: X \rightarrow P(X)$ a multivalued mapping. T is said to be a relatively KKM (R-KKM) mapping if for any $F = \{x_1, x_2, \dots, x_n\} \in \langle X \rangle$ there exists a subset $F_1 = \{x_{i_1}, x_{i_2}, \dots, x_{i_k}\}$ of F such that

$$p(e_{i_1}, e_{i_2}, \dots, e_{i_k}) \subset \bigcup_{j=1}^k Tx_{ij}$$

for a subsimplex $(e_{i_1}, e_{i_2}, \dots, e_{i_k})$ of $(e_1, e_2, \dots, e_n) = \Delta^n$ for $\{i_1, i_2, \dots, i_k\} \subset \{1, 2, \dots, n\}$.

DEFINITION 1.4 ([8]). Let $(X, H, \{p\})$ be a G - H -space, and M_1, M_2, \dots, M_n be subsets of X . A subset $F = \{x_1, x_2, \dots, x_n\}$ of n elements of X is said to be a generalized KKM (G -KKM) selection for M_1, M_2, \dots, M_n if for $\{i_1, i_2, \dots, i_k\} \subset \{1, 2, \dots, n\}$, we have

$$H(\{x_{i_1}, x_{i_2}, \dots, x_{i_k}\}) \subset \bigcup_{j=1}^k M_{ij},$$

where elements x_1, x_2, \dots, x_n are not necessarily distinct.

This generalizes the notion of a KKM selection introduced by Joo' and Kassay [4].

DEFINITION 1.5 [8]. Let $(X, H, \{p\})$ be a G - H -space, x_1, x_2, \dots, x_n be n elements of X and, M_1, M_2, \dots, M_n subsets of X . Elements x_1, x_2, \dots, x_n are called a relative KKM (R-KKM) selection for M_1, M_2, \dots, M_n if for $\{i_1, i_2, \dots, i_k\} \subset \{1, 2, \dots, n\}$, we have

$$p(e_{i_1}, e_{i_2}, \dots, e_{i_k}) \subset \bigcup_{j=1}^k M_{ij},$$

where x_1, x_2, \dots, x_n are not necessarily distinct, and $(e_{i_1}, e_{i_2}, \dots, e_{i_k})$ is a standard $n - 1$ subsimplex of (e_1, e_2, \dots, e_n) in R^n .

Clearly, every KKM selection [4] is an R-KKM selection.

Note that Definition 1.5 implies that

$$p(e_{i_1}, e_{i_2}, \dots, e_{i_k}) \subset \bigcup_{j=1}^k (M_{ij} \cap H(\{x_{i_1}, x_{i_2}, \dots, x_{i_k}\})),$$

where $\{x_{i_1}, x_{i_2}, \dots, x_{i_k}\} \subset \{x_1, x_2, \dots, x_n\}$.

Let X and Y be two sets, and $S: X \rightarrow P(Y)$ be a multivalued mapping. Then $S^{-1}: Y \rightarrow P(X)$ and $S^*: Y \rightarrow P(X)$ are defined by $x \in S^{-1}(y)$ iff $y \in Sx$ and $S^*y = X \setminus S^{-1}y$, respectively. Clearly, $x \in S^*y$ iff $y \notin Sx$.

2. R-KKM MAPPING THEOREMS

This section deals with the intersection theorems involving R-KKM mappings. The first theorem generalizes mildly [2, Theorem 1] to the case of R-KKM selections in a G - H -space.

THEOREM 2.1. Let $(X, H, \{p\})$ be a G - H -space, $F = \{x_1, x_2, \dots, x_n\} \in \langle X \rangle$, and M_1, M_2, \dots, M_n closed subsets of X . Suppose that $\{x_1, x_2, \dots, x_n\}$ is a R-KKM selection for M_1, M_2, \dots, M_n relative to a simplex $(e_1, e_2, \dots, e_n) = \Delta^n$. Then we have $\bigcap_{i=1}^n M_i \neq \emptyset$.

Proof. Since $(X, H, \{p\})$ is a G - H -space and $F = \{x_1, x_2, \dots, x_n\}$ is a R-KKM selection for the subsets M_1, M_2, \dots, M_n , we have

$$p(e_{i_1}, e_{i_2}, \dots, e_{i_k}) \subset \bigcup_{j=1}^k M_{ij} \quad \text{for } \{i_1, i_2, \dots, i_k\} \subset \{1, 2, \dots, n\},$$

where $(e_{i_1}, e_{i_2}, \dots, e_{i_k})$ is a subsimplex of $(e_1, e_2, \dots, e_n) = \Delta^n$ of R^n , and $\{x_{i_1}, x_{i_2}, \dots, x_{i_k}\} = F_1 \subset F$. If we set $E_i = p^{-1}(M_i)$ for $i = 1, 2, \dots, n$, then E_i is closed in Δ^n by the closedness of M_i and the continuity of p . Before

we can apply the KKM theorem [5], we need to show that $\text{co}(e_{i1}, e_{i2}, \dots, e_{ik}) \subset \bigcup_{j=1}^k E_{ij}$ for $\{i1, i2, \dots, ik\} \subset \{1, 2, \dots, n\}$. From the above arguments, it follows that

$$\text{co}(e_{i1}, e_{i2}, \dots, e_{ik}) \subset p^{-1} \left(\bigcup_{j=1}^k M_{ij} \right) = \bigcup_{j=1}^k (p^{-1}(M_{ij})) = \bigcup_{j=1}^k E_{ij}.$$

Hence, $\bigcap_{i=1}^n E_i \neq \emptyset$. If an element $t_0 \in \bigcap_{i=1}^n E_i$, then $t_0 \in E_i = p^{-1}(M_i)$ for all $i = 1, 2, \dots, n$. As a result, $p(t_0) \in M_i$ for all $i = 1, 2, \dots, n$, that means, $p(t_0) \in \bigcap_{i=1}^n M_i$. This completes the proof.

THEOREM 2.2. *Let X be a topological space, $(Y, H, \{p\})$ a compact G - H -space, and $T: X \rightarrow P(Y) \setminus \{\emptyset\}$ a closed valued mapping. Suppose that T is a R-KKM mapping. Then we have $\bigcap_{x \in X} Tx \neq \emptyset$.*

Proof. The proof is immediate from an application of Theorem 2.1.

THEOREM 2.3. *Let $(X, H, \{p\})$ be a compact G - H -space and $S, T: X \rightarrow P(X)$ two multivalued mappings such that:*

- (i) Tx is closed and $Sx \subset Tx$ for all $x \in X$.
- (ii) $x \in Sx$ for all $x \in X$.
- (iii) S^*x is G - H -convex for all $x \in X$.

Then $\bigcap_{x \in X} Tx \neq \emptyset$.

Proof. The proof follows from an application of Theorem 2.1 if we show that T is a R-KKM mapping, that is,

$$p(e_{i1}, e_{i2}, \dots, e_{ik}) \subset \bigcup_{j=1}^k Tx_{ij} \quad \text{for } \{i1, i2, \dots, ik\} \subset \{1, 2, \dots, n\},$$

where $(e_{i1}, e_{i2}, \dots, e_{ik})$ is a subsimplex of $(e_1, e_2, \dots, e_n) = \Delta^n$ and $F_1 = \{x_{i1}, x_{i2}, \dots, x_{ik}\}$ is a subset of $F = \{x_1, x_2, \dots, x_n\} \in \langle X \rangle$. Assume that T is not R-KKM. Then, since $(X, H, \{p\})$ is a G - H -space, we have

$$H(\{x_{i1}, x_{i2}, \dots, x_{ik}\}) \not\subset \bigcup_{j=1}^k Tx_{ij}.$$

If an element $y \in H(\{x_{i1}, x_{i2}, \dots, x_{ik}\})$, then $y \notin Tx_{ij}$ for $x_{ij} \in F_1$ for $j = 1, 2, \dots, k$. That means, $x_{ij} \in T^*y$, and as a result, $F_1 \subset T^*y$. Given that $Sx \subset Tx$ for all $x \in X$, we have $T^*y \subset S^*y$. This implies $F_1 \subset S^*y$, and by (iii), $H(F_1) \subset S^*y$. Thus, $y \in S^*y$, which implies $y \notin Sy$, a contradiction to (ii). This completes the proof.

THEOREM 2.4. *Let $(X, H, \{p\})$ be a compact G - H -space, and $S, T: X \rightarrow P(X) \setminus \{\emptyset\}$ multivalued mappings such that:*

- (i) $Sx \subset Tx$ for all $x \in X$.
- (ii) Tx is G - H -convex for all $x \in X$.
- (iii) $S^{-1}y$ is open in X for all $y \in X$.

Then there exists an element $x_0 \in X$ such that $x_0 \in Tx_0$. This is a variant form of Theorem 2.3.

3. MINIMAX INEQUALITIES

In this section, we apply the results of Section 2 to derive some minimax inequalities, which generalize some minimax results [2] in pseudoconvex spaces and H -spaces.

THEOREM 3.1. *Let $(X, H, \{p\})$ be a compact G - H -space and $f, g: X \times X \rightarrow R$ two functions such that the following assumptions hold:*

- (i) $g(x, y) \leq f(x, y)$ for all $(x, y) \in X \times X$.
- (ii) $g(x, y)$ is lower semicontinuous in its second variable, that is, the set $\{y \in X: g(x, y) \leq c \text{ for } c \in R\}$ is closed.
- (iii) $f(x, y)$ is G - H -quasiconcave in its first variable, that is, the set $\{x \in X: f(x, y) > c\}$ is G - H -convex.

Then one of the following alternatives holds:

- (A) *There exists an element y_0 of X such that $g(x, y_0) \leq c$ for all $x \in X$.*
- (B) *There is an element $x_0 \in X$ such that $f(x_0, x_0) > c$.*

COROLLARY 3.1. *Under the assumptions of Theorem 3.1, we have*

$$\inf_{y \in X} \sup_{x \in X} g(x, y) \leq \sup_{x \in X} f(x, x).$$

Proof of Theorem 3.1. Let us define two multivalued mappings S and T by $Sx = \{y \in X: f(x, y) \leq c \text{ for } c \in R\}$ and $Tx = \{y \in X: g(x, y) \leq c \text{ for } c \in R\}$, respectively. Then it follows from (i) that $Sx \subset Tx$. Consider an element $x_0 \in X$ such that $x_0 \notin Sx_0$. Then $f(x_0, x_0) > c$, which proves (B). Next, we consider the case when $x \in Sx$ for all $x \in X$. For $A \in \langle X \rangle$, $A_1 \subset A$, and $S^*y = \{x \in X: f(x, y) > c\}$, let $A_1 \subset S^*y$. Then, by (iii), it follows that $H(A_1) \subset S^*y$. Now, by Theorem 2.3, we have $\bigcap_{x \in X} Tx \neq \emptyset$, which implies (A).

REFERENCES

1. C. Bardaro and R. Ceppitelli, Some further generalizations of Knaster–Kuratowski–Mazurkiewicz theorem and minimax inequalities, *J. Math. Anal. Appl.* **132** (1988), 484–490.
2. C. Horvath, Some results on multivalued mappings and inequalities without convexity, in “Nonlinear and Convex Analysis,” (B. L. Lin and S. Simons, Eds.), Lecture Notes in Pure and Appl. Math. Series, Vol. 106, pp. 99–106, Marcel Dekker, New York, 1987.
3. I. Joo', On some convexities, *Acta Math. Hungar.* **54** (1989), 163–172.
4. I. Joo' and G. Kassay, Convexity, minimax theorems and their applications, *Ann. Univ. Sci. Budapest.* **38** (1995), 71–93.
5. B. Knaster, K. Kuratowski and S. Mazurkiewicz, Ein Beweis des fixpunktsatzes für n -dimensionale simplexe, *Fund. Math.* **14** (1929), 132–137.
6. H. Komiya, Convexity on a topological space, *Fund. Math.* **119** (1981), 107–113.
7. K. K. Tan and X. L. Zhang, Fixed point theorems on G -convex spaces and applications, *Proc. Nonlinear Funct. Anal.* **1** (1996), 1–19.
8. R. U. Verma, Generalized H-KKM mappings, *Math. Sci. Res. Hot-Line* **1** (8) (1997), 17–19.
9. R. U. Verma, Some minimax inequalities on generalized H-convex sections, *Math. Sci. Res. Hot-Line* **2** (1998), 27–32.
10. R. U. Verma, Some generalized KKM selection theorems and minimax inequalities, *Math. Sci. Res. Hot-Line* **2** (1998), 13–21.
11. R. U. Verma, RKKM mappings and their applications, *Comm. Appl. Anal.* (to appear).
12. R. U. Verma, RKKM mapping theorems and variational inequalities, *Proc. Roy. Irish Acad. Sect. A* (to appear).
13. R. U. Verma, Role of generalized KKM type selections in a class of minimax inequalities, *Appl. Math. Lett.* (to appear).